



New Jersey Center for Teaching and Learning

Progressive Science Initiative

This material is made freely available at www.njctl.org and is intended for the non-commercial use of students and teachers. These materials may not be used for any commercial purpose without the written permission of the owners. NJCTL maintains its website for the convenience of teachers who wish to make their work available to other teachers, participate in a virtual professional learning community, and/or provide access to course materials to parents, students and others.

Click to go to website:
www.njctl.org



PHYSICS

Solving Equations



April 2012

www.njctl.org

Solving for a Variable

Our goal is to be able to solve any equation for any variable that appears in it.

Let's look at a simple equation first.

$$s = \frac{d}{t}$$

The variables in this equation are s, d and t.

Solving for a variable means having it alone on the left side.

This equation is currently solved for "s".

The Rules

Like in any game there are a few rules.

1. To "undo" a mathematical operation, you must do the opposite.
2. You can do anything you want (except divide by zero) to one side of an equation, as long as you do the same thing to the other.
3. If there is more than one operation going on, you must undo them in the opposite order in which you would do them, the opposite of the "order of operations."
4. You can always switch the left and right sides of an equation.

The Rules

Let's solve this equation for "d"

$$s = \frac{d}{t}$$

That means that when we're done
we'll have d alone
on the left side of the equation.

1 First, is d already ALONE? If not, what is with it?

- A s**
- B d**
- C t**
- D it is already alone**

$$s = \frac{d}{t}$$

2 What mathematical operation connects d and t?

- A d is added to t
- B d is multiplied by t
- C d is divided by t
- D t is subtracted from d

$$s = \frac{d}{t}$$

3 What is the opposite of dividing d by t?

- A** dividing t by d
- B** dividing by s into t
- C** multiplying d by t
- D** multiplying by t by d

$$s = \frac{d}{t}$$

Rule 1. To "undo" a mathematical operation, you must do the opposite.

4 What must we also do if we multiply the right side by t?

- A divide the left side by t**
- B multiply the left side by t**
- C divide the left side by d**
- D divide the left side by d**

$$s = \frac{d}{t}$$

Rule 2. You can do anything you want (except divide by zero) to one side of an equation, as long as you do the same thing to the other.

5 Is there more than one mathematical operation acting on "d"?

- Yes**
- No**

$$s = \frac{d}{t}$$

Rule 3. If there is more than one operation going on, you must undo them in the opposite order in which you would do them, the opposite of the "order of operations."

Applying Rules 1 and 2

1. To "undo" a mathematical operation, you must do the opposite.

$$s = \frac{d}{t}$$

2. You can do anything you want (except divide by zero) to one side of an equation, as long as you do the same thing to the other. So we undo d being divided by t , by multiplying both sides by t .

$$(t) s = \frac{d}{\cancel{t}} \quad (\cancel{t})$$

$$t s = d$$

$$st = d$$

The order of the s and t doesn't matter, but we usually make them alphabetical so they look better.

Are we done?

Applying Rule 4

Rule 4. You can always switch the left and right sides of an equation.

$$\begin{array}{c} st = d \\ \swarrow \searrow \\ \boxed{d = st} \end{array}$$

We've now solved our equation for d.

A harder problem is to solve it for t.

Solving for t

Let's solve this equation for "t"

$$s = \frac{d}{t}$$

That means that when we're done
we'll have t **alone**
on the **left side** of the equation.

6 Is t already ALONE? If not, what is with it?

- A s**
- B d**
- C t**
- D it is already alone**

$$s = \frac{d}{t}$$

7 What mathematical operation connects d to t?

- A t is being divided by d**
- B d is being divided by t**
- C d is being multiplied by t**
- D t is being subtracted from d**

$$s = \frac{d}{t}$$

8 What is the opposite of dividing d by t?

- A** dividing d by t
- B** dividing s by t
- C** multiplying d by t
- D** multiplying t by d

$$s = \frac{d}{t}$$

Rule 1. To "undo" a mathematical operation, you must do the opposite.

9 What must we do if we multiply the right side by t?

- A divide the left side by t**
- B multiply the left side by t**
- C divide the left side by d**
- D divide the left side by d**

$$s = \frac{d}{t}$$

Rule 2. You can do anything you want (except divide by zero) to one side of an equation, as long as you do the same thing to the other.

10 Is there more than one mathematical operation acting on "d"?

Yes

No

$$s = \frac{d}{t}$$

Rule 3. If there is more than one operation going on, you must undo them in the opposite order in which you would do them, the opposite of the "order of operations."

Solving for T

1. To "undo" a mathematical operation, you must do the opposite.

$$s = \frac{d}{t}$$

2. You can do anything you want (except divide by zero) to one side of an equation, as long as you do the same thing to the other. So we undo d being divided by t , by multiplying both sides by t .

$$(t) s = \frac{d}{\cancel{t}} \quad (\cancel{t})$$

$$st = d$$

Are we done?

11 Is t ALONE? If not, what is with it?

- A s**
- B d**
- C t**
- D it is already alone**

st = d

12 What mathematical operation connects s to t?

- A t is being divided by d**
- B t is being divided into s**
- C t is being multiplied by s**
- D t is being subtracted from s**

$$st = d$$

13 What is the opposite of multiplying t by s?

- A dividing t by s**
- B dividing t by t**
- C multiplying t by t**
- D multiplying by t by s**

st = d

Solving for t

1. To "undo" a mathematical operation, you must do the opposite.

$$st = \frac{d}{s}$$

2. You can do anything you want (except divide by zero) to one side of an equation, as long as you do the same thing to the other.

$$\frac{\cancel{st}}{\cancel{s}} = \frac{d}{s}$$

$$t = \frac{d}{s}$$

14 Is t ALONE on the left?

- A s**
- B d**
- C t**
- D it is alone**

$$t = \frac{d}{s}$$

Solving for v_0 .

Let's solve this equation for " v_0 ."

$$v = v_0 + at$$

That means that when we're done
we'll have v_0 alone

on the left side of the equation.

15 Is v_0 already **ALONE**? If not, what is with it?

- A** only a
- B** only t
- C** at
- D** it is already alone

$$v = v_0 + at$$

16 What mathematical operation connects at to v_0 ?

- A** " at " is being divided by v_0
- B** " at " is being added to v_0
- C** v_0 is being multiplied by " at "
- D** v_0 is being divided by " at "

$$v = v_0 + at$$

17 What is the opposite of adding at to v_0 ?

- A** dividing by v_0 by "at" into t
- B** subtracting v_0 from "at"
- C** subtracting "at" from v_0
- D** dividing "at" by v_0

$$v = v_0 + at$$

18 What must we do, if we subtract "at" from the right side?

- A add "at" to the left side**
- B multiply the left side by "at"**
- C subtract "at" from the left side**
- D divide the left side by v_0**

$$v = v_0 + at$$

19 Is there more than one mathematical operation acting on "v_o"?

Yes

No

$$v = v_0 + at$$

Solving for v_0 .

1. To "undo" a mathematical operation, you must do the opposite.
2. You can do anything you want (except divide by zero) to one side of an equation, as long as you do the same thing to the other.

$$\begin{array}{r} v = v_0 + at \\ - at \quad - at \end{array}$$

$$v - at = v_0$$

$$\boxed{v_0 = v - at}$$

Solving for a

Let's solve this equation for "a"

$$v = v_0 + at$$

That means that when we're done
we'll have a alone

on the left side of the equation.

20 Is a already ALONE? If not, what is with it?

- A only v_0**
- B only t**
- C v_0 and t**
- D it is already alone**

$$v = v_0 + at$$

21 What mathematical operation connects v_0 to at ?

- A "at" is being divided by v_0**
- B v_0 is being added to "at"**
- C v_0 is being multiplied by "at"**
- D v_0 is being subtracted by "at"**

$$v = v_0 + at$$

22 What is the opposite of adding v_0 to at ?

- A** dividing by v_0 by at into t
- B** subtracting v_0 from at
- C** subtracting at from v_0
- D** dividing at by v_0

$$v = v_0 + at$$

23 What mathematical operation connects t to a?

- A a is added to t**
- B a is multiplied by t**
- C a is divided by t**
- D t is subtracted from a**

$$v = v_0 + at$$

24 What is the opposite of multiplying a by t?

- A** dividing a by t.
- B** dividing t by a.
- C** multiplying a by t
- D** multiplying t by a

$$v = v_0 + at$$

25 What must we do, if we divide by t from the right side?

- A divide the left side by a**
- B multiply the left side by a**
- C divide the left side by t**
- D multiply the left side by t**

$$v = v_0 + at$$

26 Is there more than one mathematical operation acting on "a"?

Yes

No

$$v = v_0 + at$$

27 Which operation should we undo first?

- A divide a by t**
- B subtract v_0 from at**

$$v = v_0 + at$$

28 Which operation should we undo second?

- A divide a by t**
- B subtract v_0 from at**

$$v = v_0 + at$$

Solving for a

1. To "undo" a mathematical operation, you must do the opposite.

$$\begin{array}{r} v = v_0 + at \\ - v_0 \quad - v_0 \end{array}$$

2. You can do anything you want (except divide by zero) to one side of an equation, as long as you do the same thing to the other.

$$\frac{v - v_0}{t} = \frac{at}{t}$$

3. If there is more than one operation going on, you must undo them in the opposite order in which you would do them, the opposite of the "order of operations."

$$\frac{v - v_0}{t} = a$$

Are we done?

Solving for t

Let's solve this equation for "t"

$$v = v_0 + at$$

That means that when we're done
we'll have t alone

on the left side of the equation.

Solving for t

1. To "undo" a mathematical operation, you must do the opposite.
2. You can do anything you want (except divide by zero) to one side of an equation, as long as you do the same thing to the other.
3. If there is more than one operation going on, you must undo them in the opposite order in which you would do them, the opposite of the "order of operations."

$$v = v_0 + at$$

